

# Direct resummation of the leading renormalons in inclusive semileptonic B decay

Taekoon Lee\*

*Department of Physics, Korea Advanced Institute  
of Science and Technology, Daejeon 305-701, Korea*

## Abstract

We present a Borel resummation method for the QCD corrections in inclusive, charmless, semileptonic B meson decay. The renormalon contributions are resummed to all orders by employing a bilocal expansion of the Borel transform that accurately accounts for the first infrared renormalons in the Borel plane. The renormalons in the pole mass and the QCD expansion are resummed separately, and a precise relation is obtained between a properly defined pole mass and the  $\overline{\text{MS}}$  mass. The inclusive decay rate is calculated to three loop order using an estimate of the yet unknown NNLO coefficient.

---

\*Electronic address: tlee@muon.kaist.ac.kr

The Cabibbo-Kobayashi-Maskawa matrix element  $|V_{ub}|$ , which is an important parameter in the study of CP violation in weak interactions, can be determined through the inclusive, charmless, semileptonic B decay. Presently, the largest theoretical uncertainty in the decay rate comes from the  $b$  quark mass and the QCD correction. It is well known that these two are not independent but intimately related. In the large  $b$  quark mass limit, the B meson decay rate is given by that of a free  $b$  quark with a correction quadratically suppressed in the  $b$  mass [1, 2]. The decay rate in the on-shell scheme is given in the form

$$\Gamma(B \rightarrow X_{ue\bar{\nu}_e}) = \frac{G_F^2 |V_{ub}|^2}{192\pi^3} m_b^5 f(\alpha_s) [1 + O(1/m_b^2)] \quad (1)$$

with  $m_b$  and  $f(\alpha_s)$  denoting the  $b$  quark pole mass and the QCD correction to the free  $b$  quark decay, respectively. In perturbation theory  $f(\alpha_s)$  can be expanded as

$$f = 1 + \sum_{n=0}^{\infty} f_n(\xi) \alpha_s(\mu)^{n+1}, \quad (2)$$

where  $\alpha_s(\mu)$  is the strong coupling constant and  $\xi = \mu/m_{\overline{\text{MS}}}$ , with  $m_{\overline{\text{MS}}}[\equiv m_{\overline{\text{MS}}}(m_{\overline{\text{MS}}})]$  denoting the  $\overline{\text{MS}}$  mass.

Because of the infrared (IR) renormalon the perturbative coefficients  $f_n$  are expected to grow rapidly, and this would give rise to an  $O(\Lambda_{\text{QCD}}/m_b)$  intrinsic uncertainty to the asymptotic series. On the other hand, this renormalon uncertainty in  $f(\alpha_s)$  is known to arise from the use of the pole mass in the decay rate. When the pole mass is expanded in the strong coupling it also has a renormalon caused uncertainty of  $O(\Lambda_{\text{QCD}})$ , and this uncertainty cancels that of  $f(\alpha_s)$  [3, 4]. Once the pole mass is replaced by a renormalon free mass like the  $\overline{\text{MS}}$  mass or one of the threshold masses [5, 6, 7, 8] the resulting  $f(\alpha_s)$  is in principle expected to give a well-behaved series. In practice, however, the use of the renormalon free masses are not completely satisfactory. For instance, with the  $\overline{\text{MS}}$  mass the renormalon cancellation is not so obvious at low orders and the convergence for  $f(\alpha_s)$  is quite slow, possibly due to the large power of  $m_b$  in the decay rate, and with the  $1S$  mass [7, 8] one must assume that the  $1S$  state of  $\Upsilon$  is a perturbative system.

In this paper we describe a more direct approach to the renormalon problem, in which the leading renormalons in the pole mass and  $f(\alpha_s)$  are Borel resummed separately. An essential ingredient of this program is the accurate calculation of the renormalon residue for the pole mass. The residue, which determines the normalization constant of the large order behavior, can be computed in perturbation theory [9, 10], and it turns out that the residue for the pole mass can be determined to good accuracy within a few percent error [11, 12, 13]. The renormalon cancellation then allows the residue for  $f(\alpha_s)$  to be determined to the same accuracy from the pole mass residue. The Borel transforms of the pole mass and  $f$  can be systematically expanded around the origin and the first renormalon location in the Borel plane. With the computed residues these two expansions can be interpolated to a *bilocal expansion* to obtain an accurate description of the Borel transforms in the region in-between the two expansion points. Since at the typical strong coupling in B decay the bulk of the Borel integral comes from this region of integration, the bilocal expansion can much improve the accuracy of the Bore resummation. By taking into account the renormalon singularity correctly this program essentially resums the renormalon caused large order behavior to all orders.

With the renormalon singularity on the integration contour one has to worry about the ambiguities of  $O(\Lambda_{\overline{\text{MS}}})$  for the pole mass and  $O(\Lambda_{\overline{\text{MS}}}/m_b)$  for  $f$  arising from the ambiguities

in choosing the integration contour that may either be on the upper or the lower half plane beginning at the renormalon singularities. It was shown in Refs. [14, 15] that these ambiguities cancel as long as one takes the same contour for the pole mass and  $f(\alpha_s)$ . Furthermore the authors argued that in general the pole mass, when consistently used, does not cause any ambiguity to physical observables in the heavy quark effective theory. Until recently, however, there was no procedural program implementing this observation. The main reason for this, perhaps, was that one did not know how to describe the Borel transforms beyond the immediate neighborhood of the origin in the Borel plane. Within the large  $\beta_0$  approximation [16] one can obtain the Borel transforms in the whole Borel plane, but the accuracy of the approximation is limited. What is necessary for practical applications is an accurate description of the Borel transforms in the region in-between the origin and the first renormalon singularity, since the contribution from the region far from the origin is exponentially suppressed. The bilocal expansion fills this gap.

The program described above was successfully applied to the heavy quark static potential [11] which is known to suffer from a severe convergence problem with the ordinary perturbative expansion. The Borel resummed potential obtained from employing a bilocal expansion turns out to have an excellent convergence and is in remarkable agreement with the lattice calculations.

Throughout the paper, unless stated otherwise, the perturbative expansions are assumed to be in the  $\overline{\text{MS}}$  scheme with four active quark flavors ( $N_f = 4$ ), and only the leading IR renormalon closest to the origin in the Borel plane is considered.

The pole mass in terms of the  $\overline{\text{MS}}$  mass  $m_{\overline{\text{MS}}}$  has the perturbative expansion

$$m_b = m_{\overline{\text{MS}}} \left[ 1 + \sum_{n=0}^{\infty} p_n(\xi) \alpha_s(\mu)^{n+1} \right]. \quad (3)$$

The first three coefficients  $p_n$  are known at  $\mu = m_{\overline{\text{MS}}}$  [17, 18, 19] and their  $\mu$  dependence can be obtained by the renormalization group invariance of the pole mass. The Borel resummation of this expansion can be written formally as

$$m_b[\alpha_s(\mu)] = m_{\overline{\text{MS}}} \left[ 1 + \frac{1}{\beta_0} \int_{0 \pm i\epsilon}^{\infty \pm i\epsilon} e^{-b/\beta_0 \alpha_s(\mu)} \tilde{m}(b, \xi) db \right], \quad (4)$$

where the Borel transform  $\tilde{m}(b, \xi)$  has the perturbative expansion

$$\tilde{m}(b, \xi) = \sum_{n=0}^{\infty} \frac{p_n(\xi)}{n!} \left( \frac{b}{\beta_0} \right)^n, \quad (5)$$

with  $\beta_0$  denoting the one loop coefficient of the QCD  $\beta$  function [ $\beta = d\alpha_s/d\ln(\mu^2) = -\alpha_s^2(\beta_0 + \beta_1\alpha_s + \beta_2\alpha_s^2 + \dots)$ ]. The integration contour, which can be derived from the general argument of Borel resummation of a series with a same sign large order behavior [20], can be either on the upper or the lower half plane along the positive real axis. The pole mass is known to have an ambiguity from renormalon singularity proportional to  $\Lambda_{\overline{\text{MS}}}$  [21] that has the weak coupling expansion

$$\Lambda_{\overline{\text{MS}}} \propto \mu \alpha_s(\mu)^{-\nu} e^{-\frac{1}{2\beta_0 \alpha_s(\mu)}} \left[ 1 + r_1 \alpha_s(\mu) + r_2 \alpha_s(\mu)^2 + \dots \right], \quad (6)$$

where

$$\nu = \frac{\beta_1}{2\beta_0^2}, \quad r_1 = \frac{\beta_1^2 - \beta_0 \beta_2}{2\beta_0^3},$$

$$r_2 = \frac{\beta_1^4 + 4\beta_0^3\beta_1\beta_2 - 2\beta_0\beta_1^2\beta_2 + \beta_0^2(\beta_2^2 - 2\beta_1^3) - 2\beta_3\beta_0^4}{8\beta_0^6}. \quad (7)$$

For the Borel integral (4) to generate this ambiguity one can easily see that the Borel transform  $\tilde{m}(b, \xi)$  must have the singularity

$$\tilde{m}(b, \xi) = \frac{C_m \xi}{(1-2b)^{1+\nu}} [1 + c_1(1-2b) + c_2(1-2b)^2 + \dots] + \text{analytic part}, \quad (8)$$

where

$$c_1 = \frac{r_1}{2\nu\beta_0}, \quad c_2 = \frac{r_2}{4\nu(\nu-1)\beta_0^2}. \quad (9)$$

The ‘analytic part’ denotes any terms analytic around  $b = 1/2$  with radius of convergence bounded by the next renormalon at  $b = 3/2$  and  $C_m$  is a real constant.

With this singularity in  $\tilde{m}(b, \xi)$  the resummed pole mass can be written formally as

$$m_b = m_{\text{BR}}[\alpha_s(\mu)] \pm i\Gamma_m[\alpha_s(\mu)], \quad (10)$$

where the ‘Borel resummed (BR)’ mass  $m_{\text{BR}}$  denotes the real part of the Borel integral in Eq. (4) and  $\Gamma_m$  denotes the imaginary part obtained with the integration contour on the upper half plane. Note that by definition the BR mass has the same perturbative expansion as the pole mass and is not a short distance mass.

The residue  $C_m$ , which is not known exactly, can be calculated systematically in perturbation theory [9, 10]. The details of the calculation adopted here can be found in [11]. To compute the residue we first expand using the known  $p_n$  up to NNLO and the unknown  $p_3$  the function  $R(b, \xi) \equiv (1-2b)^{(1+\nu)}\tilde{m}(b, \xi)$  about the origin in the  $w$  plane defined by the following conformal mapping

$$w = \frac{\sqrt{1+b} - \sqrt{1-2b/3}}{\sqrt{1+b} + \sqrt{1-2b/3}}, \quad (11)$$

which gives to  $O(w^3)$

$$R[b(w), 1] = 0.42441 + 0.61198w + 0.25351w^2 + [-108.99217 + 7.90052p_3(1)]w^3. \quad (12)$$

This allows us to estimate  $p_3$  [22]. From the pattern of the known lower order terms we may safely assume  $|108.99217 - 7.90052p_3(1)| < 1$ , which leads to

$$p_3(1) = 13.796 \pm 0.127. \quad (13)$$

Now evaluating Eq. (12) at the renormalon location  $w=1/5$  we obtain the residue

$$C_m = R[b(w=1/5), 1] \approx 0.42441 + 0.12240 + 0.01014 \pm 0.00800 = 0.55695 \pm 0.00800, \quad (14)$$

where the error comes from the uncertainty in the estimate (13).

The BR mass  $m_{\text{BR}}$  can be obtained accurately by an interpolation of the two expansions (5) and (8) of the Borel transform  $\tilde{m}(b, \xi)$  (bilocal expansion [11]). Since at the typical values of the strong coupling considered here the bulk of the contribution to the Borel integral comes from the region between the origin and the first IR renormalon, it is important to have the Borel transform in this region as accurately as possible from the known first perturbative

terms. As demonstrated in the heavy quark potential the accuracy of the Borel transform can be improved greatly through the bilocal expansion [11].

The bilocal expansion of the Borel transform for the pole mass is given by

$$\begin{aligned}\tilde{m}(b, \xi) &= \lim_{N, M \rightarrow \infty} \tilde{m}_{N, M}(b, \xi) \\ &= \lim_{N, M \rightarrow \infty} \left\{ \sum_{n=0}^N \frac{q_n(\xi)}{n!} \left( \frac{b}{\beta_0} \right)^n + \frac{C_m \xi}{(1-2b)^{1+\nu}} \left[ 1 + \sum_{i=1}^M c_i (1-2b)^i \right] \right\},\end{aligned}\quad (15)$$

where  $q_n$  are to be determined by demanding Eq. (15) reproduce Eq. (5) when expanded about  $b = 0$ . With the known first two coefficients  $c_{1,2}$ , and taking  $M = 2$ , we have the first three  $q_n$  as

$$\begin{aligned}q_0(\xi) &= p_0 - C_m \xi (1 + c_1 + c_2), \\ q_1(\xi) &= p_1(\xi) - 2C_m \xi \beta_0 [1 - c_2 + \nu(1 + c_1 + c_2)], \\ q_2(\xi) &= p_2(\xi) - 4C_m \xi \beta_0^2 [2 + \nu(3 + c_1 - c_2) + \nu^2(1 + c_1 + c_2)].\end{aligned}\quad (16)$$

Since the two expansions about the origin and about the renormalon singularity are expected to be convergent on the disks,  $|b| < 1/2$  and  $|b - 1/2| < 1$ , respectively, the interpolation (15) is expected to give a good description of the Borel transform in the region that includes the two expansion points.

Taking the canonical value for the strong coupling  $\alpha_s(m_{\overline{\text{MS}}}) = 0.22$  and  $\xi = 1$ , and substituting the known  $\tilde{m}_{N,2}(b, 1)$ , ( $N = 0, 1, 2$ ), into the Borel integral (4), and performing the integration in the  $w$ -plane for the convenience of numerical integration, we obtain

$$m_{\text{BR}} = m_{\overline{\text{MS}}}(1 + 0.15769 + 0.00409 - 0.00028 \mp 0.00014),\quad (17)$$

where the error was obtained by varying  $C_m$  within the uncertainty in the computed residue in Eq. (14). Notice the remarkable convergence of the BR mass compared to the power series (3) which at  $\xi = 1$  gives  $m_b = m_{\overline{\text{MS}}}(1 + 0.09337 + 0.04550 + 0.03235)$ . This shows that the bilocal expansion improves the convergence by two orders of magnitude.

We now turn to the resummation of  $f(\alpha_s)$ . For this we first note that because of the large power ( $m_b^5$ ) in the pole mass in the decay rate the expansion (2) may not be an optimal expansion for  $f(\alpha_s)$ . We can reorganize the expansion (2) by writing

$$f[\alpha_s(\mu)] = F(1 + H[\alpha_s(\mu)])\quad (18)$$

where  $F(x)$  is an arbitrary function analytic at  $x = 1$ . Since  $H$  itself can be regarded as a coupling constant, with its relation to the  $\overline{\text{MS}}$  coupling constant defined implicitly by Eq. (18), we see that a particular choice of  $F$  is nothing but a renormalization scheme choice.

Given an  $F$  the perturbative expansion for  $H$

$$H = \sum_{n=0}^{\infty} h_n(\xi) \alpha_s(\mu)^{n+1}\quad (19)$$

can be easily obtained using the definition (18) and the series (2) for  $f(\alpha_s)$ . Instead of resumming the series (2) directly we shall resum the expansion (19) for  $H$ .

The Borel resummation for  $H$  can be done in a similar fashion as in the pole mass resummation, assuming that the first IR renormalon for  $H$  is known. Then with the resummed pole mass and  $H$  we have

$$m_b^5 f = (m_{\text{BR}} \pm i\Gamma_m)^5 F(1 + H_{\text{BR}} \pm i\Gamma_H),\quad (20)$$

where  $H_{\text{BR}}$  denotes the real part of the Borel integral

$$H[\alpha_s(\mu)] = \frac{1}{\beta_0} \int_{0 \pm i\epsilon}^{\infty \pm i\epsilon} e^{-b/\beta_0 \alpha_s(\mu)} \tilde{H}(b, \xi) db \quad (21)$$

and  $\Gamma_H$  denotes the imaginary part obtained with the contour on the upper half plane. The Borel transform  $\tilde{H}(b, \xi)$  has the perturbative expansion

$$\tilde{H}(b, \xi) = \sum_{n=0}^{\infty} \frac{h_n(\xi)}{n!} \left( \frac{b}{\beta_0} \right)^n. \quad (22)$$

The structure of the IR renormalon for  $H$  can be easily obtained by the cancellation of the renormalons in  $m_b^5$  and  $f(\alpha_s)$ . Expanding Eq. (20) in powers of  $\Gamma_{m,H}$  we obtain

$$m_b^5 f = m_{\text{BR}}^5 F(1 + H_{\text{BR}}) \pm i m_{\text{BR}}^4 [5F(1 + H_{\text{BR}})\Gamma_m + F'(1 + H_{\text{BR}})m_{\text{BR}}\Gamma_H] + O(\Gamma_{m,H}^2) \quad (23)$$

The cancellation of the renormalons means vanishing of the ambiguous imaginary part, which gives

$$\Gamma_H(\alpha_s) = -\frac{5F[1 + H_{\text{BR}}(\alpha_s)]}{F'[1 + H_{\text{BR}}(\alpha_s)]m_{\text{BR}}(\alpha_s)}\Gamma_m(\alpha_s). \quad (24)$$

The renormalon singularity for  $\tilde{H}(b, \xi)$  can now be easily obtained by expanding the right-hand-side of this equation in small  $\alpha_s$ .

At this point we choose a specific form for  $F(x)$ . Considering the large power of the pole mass in the decay rate a plausible choice for  $F$  is

$$F(x) = \left[ 1 + \frac{1}{q}(x - 1) \right]^q, \quad (25)$$

with  $q$  a number not necessarily an integer.  $F(1 + H)$  is then essentially a power function with a rescaling of the normalization for  $H$  that is for the purpose of convenience only and does not have any physical effect.

Using the perturbative expansions in Eqs. (3) and (19) of  $m_{\text{BR}}$  and  $H_{\text{BR}}$  (which have the identical expansions with  $m_b$  and  $H$ , respectively) and  $\Gamma_m$  obtained from the renormalon singularity (8), we have

$$\Gamma_H(\alpha_s) = \frac{5C_m \xi \Gamma(-\nu) \sin(\nu\pi)}{(2\beta_0)^{1+\nu}} \alpha_s^{-\nu} e^{-\frac{1}{2\beta_0 \alpha_s}} \left[ 1 + \tilde{r}_1 \alpha_s + \tilde{r}_2(\xi) \alpha_s^2 + \dots \right], \quad (26)$$

where

$$\begin{aligned} \tilde{r}_1 &= r_1 - p_0 + h_0/q \\ \tilde{r}_2(\xi) &= r_2 - p_1(\xi) + p_0(p_0 - r_1) + [h_1(\xi) + h_0 r_1 - p_0 h_0]/q, \end{aligned} \quad (27)$$

with  $r_{1,2}$  given in Eq. (7). For the Borel integral (21) to have this imaginary part the Borel transform  $\tilde{H}(b, \xi)$  must have the singularity

$$\tilde{H}(b, \xi) = \frac{C_H(\xi)}{(1 - 2b)^{1+\nu}} \left[ 1 + \tilde{c}_1(1 - 2b) + \tilde{c}_2(\xi)(1 - 2b)^2 + \dots \right] + \text{analytic part}, \quad (28)$$




FIG. 1: The ratio of the computed renormalon residue to the expected one from the renormalon cancellation at NLO (solid) and at estimated NNLO (dashed) vs renormalization scale  $\xi = \mu/m_{\overline{\text{MS}}}$ .

with

$$C_H(\xi) = -5C_m\xi, \quad \tilde{c}_1 = \frac{\tilde{r}_1}{2\nu\beta_0}, \quad \tilde{c}_2(\xi) = \frac{\tilde{r}_2(\xi)}{4\nu(\nu-1)\beta_0^2}. \quad (29)$$

Now the decay rate in the BR scheme is given by

$$\Gamma(b \rightarrow X_{ue\bar{\nu}_e}) = \frac{G_F^2 |V_{ub}|^2}{192\pi^3} m_{\text{BR}}^5 f_{\text{BR}}(\alpha_s) [1 + O(1/m_{\text{BR}}^2)], \quad (30)$$

where  $f_{\text{BR}} = F(1 + H_{\text{BR}})$ . Note that this is nothing but the decay rate in the on-shell scheme with Borel resummation, but now there is no theoretical ambiguity inherent in the formula in Eq. (1). Unlike  $f_{\text{BR}}$ , for instance,  $f(\alpha_s)$  in Eq. (1) is not well defined due to the divergence of the expansion (2).

We now take the  $q \rightarrow \infty$  limit in Eqs. (25) and (27), which is equivalent to putting  $f = \exp(H)$ . What is nice with this limit is that the renormalon cancellation now occurs within the sum of the series for  $H$  and that of  $5\ln(m_b)$ . We think this is an attractive feature, considering the good cancellation of renormalons within the sum of the pole mass and the static potential in heavy quarkonium system [12, 13]. Also, we shall see that the  $q$  dependence of  $f(\alpha_s)$  becomes minimal in this limit.

Before starting computation of  $H_{\text{BR}}$  we first check how well the renormalon residue  $C_H(\xi)$  computed using the known NLO calculation for  $H$  compares to the one in Eq. (29) expected from the renormalon cancellation. This exercise will allow us to estimate the unknown NNLO coefficient for  $f$ . The calculation for the residue  $C_H(\xi)$  is completely similar to that of the pole mass residue  $C_m$ . The ratio of the calculated NLO residue to the expected residue is plotted in Fig. 1. The best matching between the calculated and the expected occurs at  $\xi \approx 0.34$  with about 80% agreement. This scale is smaller than  $b$  quark mass, and this is in line with the expectation that the optimal expansion for  $f(\alpha_s)$  would occur at a scale somewhat below the  $b$  mass. This also shows that the NNLO contribution to the residue should be significant. Now the unknown NNLO coefficient  $f_2$  can be estimated in the following way. We demand the NNLO residue at  $\xi=0.34$  fall within a conservative 20% of the expected value ( $|\text{ratio} - 1| \leq 0.2$ ), which gives

$$0.8 \leq -1.552 - 0.277 f_2(1) \leq 1.2. \quad (31)$$




FIG. 2:  $f_{\text{BR}}$  at LO (dotted), NLO (dashed), and estimated NNLO (solid) at  $\alpha_s(m_{\overline{\text{MS}}}) = 0.22$  vs renormalization scale  $\xi = \mu/m_{\overline{\text{MS}}}$ . The shaded band shows the uncertainty of  $f_{\text{BR}}$  at NNLO.

This provides us an estimate of the NNLO coefficient:

$$f_2(1) = -9.22 \pm 0.72. \quad (32)$$

This corresponds to  $f = [1 - 2.41\bar{a}_s - 21.30\bar{a}_s^2 - (286 \pm 22)\bar{a}_s^3]$  and  $m_b^5 f = m_{\overline{\text{MS}}}^5 [1 + 4.25\bar{a}_s + 26.78\bar{a}_s^2 + (160 \pm 22)\bar{a}_s^3]$  where  $\bar{a}_s = \alpha_s(m_{\overline{\text{MS}}})/\pi$ . The latter shows that our estimate of the NNLO coefficient sits in the middle of the estimate  $5^3 = 125$  based on a naive power growing behavior [23] and the Padé estimate 188 [24].

The computation of  $H_{\text{BR}}$  is completely parallel to that of  $m_{\text{BR}}$ . We first combine the two expansions in Eqs. (22) and (28) for  $H$  into the bilocal expansion  $\tilde{H}_{N,M}(b, \xi)$  defined similarly to that of the pole mass in Eq. (15). The first two  $\tilde{H}_{N,2}(b, \xi)$  ( $N = 0, 1$ ) can be obtained from the known NLO expansions [25] of  $f(\alpha_s)$ , and the NNLO  $\tilde{H}_{2,2}(b, \xi)$  from the estimated coefficient in Eq. (32). Substituting these into the Borel integral (21) and performing the integration the Borel resummed  $H_{\text{BR}}$  up to NNLO is obtained. The result for  $f_{\text{BR}} = \exp(H_{\text{BR}})$  is in Fig. 2. The plots show that the scale dependence is significant at LO but becomes milder at NLO and small at NNLO for  $\xi \geq 0.5$ . Note that the NLO result is completely within the NNLO band arising from the error in the  $f_2$  estimate (32). From the perspective of scale dependence the estimate in Eq. (32) with the lower bound in the error estimate is preferred because it defines the lower bound of the NNLO band which has a less scale dependence than the upper one. In the presence of this unphysical scale dependence we can take the principle of minimal sensitivity (PMS) [26] values as the predictions of the NNLO resummations. This gives

$$f_{\text{BR}} = e^{H_{\text{BR}}} = 0.728 \pm 0.011, \quad (33)$$

where the error was taken from the difference between the PMS values of the central line and the lower bound of the NNLO band. Since an error estimate is always subjective, a more conservative estimate can be obtained, for example, by taking the variation of the NNLO estimates at  $\xi = 0.4$  as the uncertainty, which gives  $f_{\text{BR}} \approx 0.728 \pm 0.017$ .

We now come back to the case of a finite value of  $q$ . In principle  $f_{\text{BR}}(\alpha_s)$  should not depend on  $q$ , but with a finite order perturbation there is always scheme dependence, and a






FIG. 3: Renormalization scheme dependence of  $f_{\text{BR}}$  at NNLO (estimated).

scheme with a less scheme dependence is obviously preferable. To see the  $q$  dependence we repeated the NNLO computation of  $f_{\text{BR}}$  at various values of  $q$ , and the result is in Fig. 3. The scheme dependence at small  $q$  is quite significant, for example, the difference between  $q = 1$  and  $q = 5$  is larger than the NNLO contribution at  $\xi \geq 0.25$ . For large  $q$  the scheme dependence is very mild and becomes minimal at  $q = \infty$ . This shows that  $q = 1$ , the usual choice in the literature, is not a good scheme, but rather  $q = \infty$  is a good scheme to choose.

Combining  $m_{\text{BR}}$  and  $f_{\text{BR}}$  in Eqs. (17) and (33), respectively, we finally obtain the B decay rate given in Eq. (30) with

$$m_{\text{BR}}^5 f_{\text{BR}} = m_{\overline{\text{MS}}}^5 (1.539 \pm 0.023). \quad (34)$$

To conclude, we summarize our results as follows.

- The leading renormalon in the pole mass can be resummed accurately by employing a bilocal expansion. The BR mass  $m_{\text{BR}}$ , which is the real part of the integral (4), is unambiguous and converges rapidly, although it has the same divergent perturbative expansion as the pole mass and is not a short distance mass. The renormalon ambiguities in the pole mass cancel when a consistent integration contour is taken for all Borel integrals. Defined as the pole mass, the BR mass gives a precise meaning to the relation between the pole mass and a high energy mass like the  $\overline{\text{MS}}$  mass.
- A leading renormalon in the effective field theories is no longer a huddle but a blessing for the perturbation theory. By providing a manageable constraint on the functional form of the Borel transform it allows one to speed up the convergence of perturbation theory.
- The renormalon residue of the pole mass can be determined accurately. Using the renormalon cancellation one can determine the residue of the other perturbative expansion that cancels the renormalon ambiguity in the pole mass to the same accuracy. In the case of the B decay, we used this information on the residue to estimate the NNLO coefficients  $f_2$  in Eq. (32).
- The renormalization scheme dependence of  $f(\alpha_s)$ , parametrized by Eq. (25), is significant and should be taken into account.

- The B decay rate was calculated to NNLO (using an estimate of  $f_2$ ) in the BR scheme.

## Acknowledgments

This work was supported in part by BK21 Core Project.

- 
- [1] J. Chay, H. Georgi, and B. Grinstein, Phys. Lett. **B247**, 399 (1990).
  - [2] I. I. Y. Bigi, N. G. Uraltsev, and A. I. Vainshtein, Phys. Lett. **B293**, 430 (1992), hep-ph/9207214.
  - [3] I. I. Y. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D50**, 2234 (1994), hep-ph/9402360.
  - [4] M. Beneke, V. M. Braun, and V. I. Zakharov, Phys. Rev. Lett. **73**, 3058 (1994), hep-ph/9405304.
  - [5] I. I. Y. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D52**, 196 (1995), hep-ph/9405410.
  - [6] M. Beneke, Phys. Lett. **B434**, 115 (1998), hep-ph/9804241.
  - [7] A. H. Hoang and T. Teubner, Phys. Rev. **D60**, 114027 (1999), hep-ph/9904468.
  - [8] A. H. Hoang, Z. Ligeti, and A. V. Manohar, Phys. Rev. Lett. **82**, 277 (1999), hep-ph/9809423.
  - [9] T. Lee, Phys. Rev. **D56**, 1091 (1997), hep-th/9611010.
  - [10] T. Lee, Phys. Lett. **B462**, 1 (1999), hep-ph/9908225.
  - [11] T. Lee Phys. Rev. **D67**, 014020 (2003), hep-ph/0210032.
  - [12] A. Pineda, JHEP **06**, 022 (2001), hep-ph/0105008.
  - [13] A. Pineda (2002), hep-ph/0208031.
  - [14] M. E. Luke, A. V. Manohar, and M. J. Savage, Phys. Rev. **D51**, 4924 (1995), hep-ph/9407407.
  - [15] M. Neubert and C. T. Sachrajda, Nucl. Phys. **B438**, 235 (1995), hep-ph/9407394.
  - [16] M. Beneke and V. M. Braun, Phys. Lett. **B348**, 513 (1995), hep-ph/9411229.
  - [17] N. Gray, D. J. Broadhurst, W. Grafe, and K. Schilcher, Z. Phys. **C48**, 673 (1990).
  - [18] K. Melnikov and T. v. Ritbergen, Phys. Lett. **B482**, 99 (2000), hep-ph/9912391.
  - [19] K. G. Chetyrkin and M. Steinhauser, Nucl. Phys. **B573**, 617 (2000), hep-ph/9911434.
  - [20] T. Lee, Phys. Rev. **D66**, 034027 (2002), hep-ph/0104306.
  - [21] M. Beneke, Phys. Lett. **B344**, 341 (1995), hep-ph/9408380.
  - [22] K. S. Jeong and T. Lee, Phys. Lett. **B550**, 166 (2002), hep-ph/0204150.
  - [23] I. I. Y. Bigi, M. A. Shifman, N. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D56**, 4017 (1997), hep-ph/9704245.
  - [24] M. R. Ahmady, F. A. Chishtie, V. Elias, and T. G. Steele, Phys. Lett. **B479**, 201 (2000), hep-ph/9910551.
  - [25] T. van Ritbergen, Phys. Lett. **B454**, 353 (1999), hep-ph/9903226.
  - [26] P. M. Stevenson, Phys. Rev. **D23**, 2916 (1981).